

DBCg – SKAGEN TRIAL 1

DANISH BREAST CANCER COOPERATIVE GROUP

COSMETIC SCORING FORM

0-10 years

Name – Patient ID				Hospital						
<div style="display: flex; justify-content: space-around; align-items: flex-end;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> - <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> - <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> - <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div> <div style="display: flex; justify-content: space-around; font-size: small; margin-top: 2px;"> Day Month Year No. </div>										
Start year	<input style="width: 50px; height: 20px;" type="text"/>	Years after RT (0 is before RT)		0	1	2	3	4	5	10
		Day, month								
Oncoplastic surgery before RT	0 = No 1 = Yes									
Weight (kg)										
Height (cm)										
Tobacco: Package years										
Tobacco: Current smoker	0 = No 1 = Yes									
Specialist evaluated morbidity										
Breast reconstruction after mastectomy	0 = No 1 = Yes 1)									
Dyspigmentation, Breast / Chest wall	2)									
Dyspigmentation, Boost	2) 4 = Boost not given									
Telangiectasiae, Breast / Chest wall	0 = None, 1 = <1/cm ² , 2 = 1-4/cm ² 3 = >4/cm ²									
Telangiectasiae, Boost	0 = None, 1 = <1/cm ² , 2 = 1-4/cm ² , 3 = >4/cm ² , 4 = Boost not given									
Fibrosis, Breast / Chest wall	0 = None, 1 = Slightly palpable, 2 = Palpable 3 = Clearly palpable, retraction of skin and fixation									
Fibrosis, Boost	0 = None, 1 = Slightly palpable, 2 = Palpable, 3 = Clearly palpable, retraction of skin and fixation, 4 = Boost not given									
Scar	3)									
Edema, Breast / Chest wall	0 = None, 1 = Asymptomatic, 2 = Symptomatic, 3 = Secondary dysfunction									
Total cosmetic assessment (subjective) (After breast conserving surgery only.)	4)									
Patient wishes to leave the RT protocol	0 = No 1 = Yes									
Other e.g. postoperative seromen / haematoma or morbidity of heart / lung / thorax wall:										
1) If yes, do not fill out: Dyspigmentation, Telangiectasiae, Fibrosis, Edema, Scar, Total cosmetic assessment (subjective)										
2) 0 = No difference in color both for skin and papilla. 1 = Papilla <u>or</u> skin lighter/darker than on non-irradiated breast. 2 = <u>Both</u> papilla <u>and</u> skin lighter/darker than on non-irradiated breast. (Not applicable after mastectomy.) 3 = Dramatic difference in color between irradiated and non-irradiated breast, either on the papilla and/or on skin.										
3) 0 = Largely not visible. 1 = Visible but does not influence the cosmetic result. 2 = Visible and affects the cosmetic result to some degree. 3 = Visible and affects the cosmetic result considerably, 4 = not applicable.										
4) 0 = Excellent. No asymmetry, normal contour without visible deformities or skin changes and only slight increase in consistency. 1 = Good. Slight asymmetry and/or slight deformity and/or slightly increased pigmentation / telangiectasia and/or slight edema of breast and/or some increase in breast consistency. 2 = Fair. Distinct asymmetry and/or distinct deformity and/or distinct increased pigmentation / telangiectasia and/or edema of the breast and/or scattered fibrosis. 3 = Poor. Considerable asymmetry and/or severe deformity and/or considerable dyspigmentation / telangiectasia and/or edema of the breast and/or severe fibrosis or necrosis.										

Patient ID - - -

Day Month Year No.

Years after RT (0 is before RT)	0	1	2	3	4	5	10
---------------------------------	---	---	---	---	---	---	----


Measurement of arm lymphedema

Does the patient have an arm sleeve?	0 = No 1 = Yes								
If yes, how often is the arm sleeve used?	5)								
Has the sleeve been in use within the last 24 hours?	0 = No 1 = Yes								
Does the patient have a compression glove?	0 = No 1 = Yes								
If yes, how often is the compression glove used?	5)								
Measurement of arm (mm), 15 cm proximal of the olecranon 6)	Treated side								
	Opposite side								
Measurement of arm (mm), 10 cm distal of the olecranon 6)	Treated side								
	Opposite side								
Range of shoulder motion (degrees), Abduction 7)	Treated side								
	Opposite side								
Range of shoulder motion (degrees), Flexion 7)	Treated side								
	Opposite side								
Has the patient had a skin infection in the breast / arm on the treated side (before RT / since last follow up)?	0 = No 1 = Yes								


5) 4 = Every day or almost every day.
3 = Every week.
2 = 1-3 times per month.
1 = More seldom.

6) **Definition of edema:** The absolute measurements on the arms are reported (thus no subtraction is done in the reporting of data).

7) **Definition of ROM:** Range of shoulder motion is measured at abduction and flexion bilaterally. An imaginary line in the middle of the upper arm is projected on the circular scale, and the number of degrees is reported. Absolute values are reported separately for each arm.



Abduction



Flexion