



Name, personal identifier		Epirubicin (mg/m <sup>2</sup> )		Starting dose		Dose-Level -1		Dose-Level -2		Dose Level -3		
				90		75		60		Discontinue		
		Cyclophosphamide (mg/m <sup>2</sup> )		600		500		400		Discontinue		
Height (cm):	Weight (kg):	Surface (m <sup>2</sup> ):		Reduce starting dose-level in the presence of severe co-morbidity								
Year <input type="text"/>	Week no.	0	1		2	3		4	5		6	7
	Day, month											
		1. cycle		2. cycle		3. cycle		4. cycle				
Epirubicin (mg)												
Cyclophosphamide (mg)												
Growth factor support		0=no, 1=yes										
Pembrolizumab (mg)												
Ovarian function suppression		0=no, 1=yes										
Zoledronic acid (mg)												
<b>Hematological toxicity</b> If absolute neutrophil count (ANC) < 1.0 x 10 <sup>9</sup> /L on a day of planned treatment postpone treatment until ANC ≥1.0 and if ANC 0.5-0.9 on day one keep starting dose. On the second occasion reduce by one level. If ANC<0.5 postpone until ANC≥1.0 and reduce by one dose level. If platelets < 100 x 10 <sup>9</sup> /L on a day of planned treatment postpone treatment until platelets ≥100 and keep dose level if platelets 50-99 (reduce by one dose level on second occasion) and if platelets <50 reduce by one dose level.												
Enter on day 1	Neutrophils (x10 <sup>9</sup> /L)											
	Leucocytes (x10 <sup>9</sup> /L)											
	Haemoglobine (mmol/L)											
	Thrombocytes (x10 <sup>9</sup> /L)											
<b>Non-hematological toxicity</b> Grade 3 and 4 should result in dose reduction												
Febrile neutropenia		0 – 4										
Mouth or throat sores		0 – 2										
Nausea		0 – 2										
Vomiting		0 – 4										
Diarrhea		0 – 4										
Rash		0 – 2										
Hand-foot syndrome		0 – 2										
Myalgia		0 – 2										
Arthralgia		0 – 2										
Numbness & tingling		0 – 4										
Cooling gloves used		0=no, 1=yes										
Fatigue		0 – 2										
Missed expected menses		0=no, 1=yes										
Other, add text		0 – 4										
Other, add text		0 – 4										
In case of any grade 3 or 4 toxicity make a record in the study specific CRF or eJournal of day of start and resolution as well as relation to the study drugs (related, possible related, or not related).												



Name, personal identifier	Paclitaxel (mg/m <sup>2</sup> ), A +C Carboplatin (AUC), C	Starting dose	Dose-Level -1	Dose-Level -2	Dose Level -3
		80	65	50	Discontinue
		5	4	3	Discontinue

Height (cm):	Weight (kg):	Surface (m <sup>2</sup> ):	Reduce starting dose-level in the presence of severe co-morbidity
--------------	--------------	----------------------------	---

Year <input type="text"/>	Week no.	9	10	11	12	13	14	15	16	17	18	19	20
	Day, month												
		5. cycle			6. cycle			7. cycle			8. cycle		
Paclitaxel (mg), iv. day 1+8+15													
Carboplatin (mg), iv. day 1													
Growth factor support 0=no, 1=yes													
Pembrolizumab (mg)													
Ovarian function suppression 0=no, 1=yes													
Zoledronic acid (mg)													

**Hematological toxicity** If absolute neutrophil count (ANC) < 1.0 x 10<sup>9</sup>/L on a day of planned treatment postpone treatment until ANC ≥1.0 and **add** growth factor support. On the second occasion reduce by one level. If platelets < 100 x 10<sup>9</sup>/L on a day of planned treatment postpone treatment until platelets ≥100 and reduce by one dose level.

Enter on day 1	Neutrophils (x10 <sup>9</sup> /L)												
	Leucocytes (x10 <sup>9</sup> /L)												
	Hemoglobin (mmol/L)												
	Thrombocytes (x10 <sup>9</sup> /L)												

**Renal function**

On day 1	GFR ml/min												
----------	------------	--	--	--	--	--	--	--	--	--	--	--	--

**Non-hematological toxicity** Grade 3 and 4 should result in dose reduction

Febrile neutropenia	0 – 4												
Mouth or throat sores	0 – 2												
Nausea	0 – 2												
Vomiting	0 – 4												
Diarrhea	0 – 4												
Rash	0 – 2												
Hand-foot syndrome	0 – 2												
Myalgia	0 – 2												
Arthralgia	0 – 2												
Numbness & tingling	0 – 4												
Cooling gloves used	0=no, 1=yes												
Fatigue	0 – 2												
Missed expected menses	0=no, 1=yes												
Other, add text	0 – 4												
Other, add text	0 – 4												

In case of any grade 3 or 4 toxicity make a record in the study specific CRF or eJournal of day of start and resolution as well as relation to the study drugs (related, possible related, or not related).