



| | | | | | | | | | | | | | | |
|---|------------------------------------|---------------------------------------|---|----------------------------|----------|---|---|---------------|---|---------------|----------|---|----|----|
| Name, personal identifier | | Epirubicin (mg/m ²) | | Starting dose | | Dose-Level -1 | | Dose-Level -2 | | Dose Level -3 | | | | |
| | | | | 75 | | 60 | | 50 | | Discontinue | | | | |
| | | | | 600 | | 500 | | 400 | | Discontinue | | | | |
| | | Cyclophosphamide (mg/m ²) | | 900 x 2 | | 700 x 2 | | 500 x 2 | | 300 x 2 | | | | |
| Capecitabine (mg/m ²) | | | | | | | | | | | | | | |
| Hight (cm): | | Weight (kg): | | Surface (m ²): | | Reduce starting dose-level in the presence of severe co-morbidity | | | | | | | | |
| Year <input type="text"/> | Week no. | | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| | Day, month | | | | | | | | | | | | | |
| | | 1. cycle | | | 2. cycle | | | 3. cycle | | | 4. cycle | | | |
| Epirubicin (mg) | | | | | | | | | | | | | | |
| Cyclophosphamide (mg) | | | | | | | | | | | | | | |
| Capecitabine (mg days 1-14) | | | | | | | | | | | | | | |
| Growth factor support | | 0=no, 1=yes | | | | | | | | | | | | |
| Pembrolizumab (mg) | | | | | | | | | | | | | | |
| Ovarian function suppression | | 0=no, 1=yes | | | | | | | | | | | | |
| Zoledronic acid (mg) | | | | | | | | | | | | | | |
| Hematological toxicity If absolute neutrophil count (ANC) < 1.0 x 10 ⁹ /L on a day of planned treatment postpone treatment until ANC ≥1.0 and if ANC 0.5-0.9 on day one keep starting dose. On the second occasion reduce by one level. If ANC<0.5 postpone until ANC≥1.0 and reduce by one dose level. If platelets < 100 x 10 ⁹ /L on a day of planned treatment postpone treatment until platelets ≥100 and keep dose level if platelets 50-99 (reduce by one dose level on second occasion) and if platelets <50 reduce by one dose level. | | | | | | | | | | | | | | |
| Enter on day 1 | Neutrophils (x10 ⁹ /L) | | | | | | | | | | | | | |
| | Leucocytes (x10 ⁹ /L) | | | | | | | | | | | | | |
| | Haemoglobine (mmol/L) | | | | | | | | | | | | | |
| | Thrombocytes (x10 ⁹ /L) | | | | | | | | | | | | | |
| Non-hematological toxicity Grade 3 and 4 should result in dose reduction | | | | | | | | | | | | | | |
| Febrile neutropenia | | 0 – 4 | | | | | | | | | | | | |
| Mouth or throat sores | | 0 – 2 | | | | | | | | | | | | |
| Nausea | | 0 – 2 | | | | | | | | | | | | |
| Vomiting | | 0 – 4 | | | | | | | | | | | | |
| Diarrhea | | 0 – 4 | | | | | | | | | | | | |
| Rash | | 0 – 2 | | | | | | | | | | | | |
| Hand-foot syndrome | | 0 – 2 | | | | | | | | | | | | |
| Myalgia | | 0 – 2 | | | | | | | | | | | | |
| Arthralgia | | 0 – 2 | | | | | | | | | | | | |
| Numbness & tingling | | 0 – 4 | | | | | | | | | | | | |
| Cooling gloves used | | 0=no, 1=yes | | | | | | | | | | | | |
| Fatigue | | 0 – 2 | | | | | | | | | | | | |
| Missed expected menses | | 0=no, 1=yes | | | | | | | | | | | | |
| Other, add text | | 0 – 4 | | | | | | | | | | | | |
| Other, add text | | 0 – 4 | | | | | | | | | | | | |
| In case of any grade 3 or 4 toxicity make a record in the study specific CRF or eJournal of day of start and resolution as well as relation to the study drugs (related, possible related, or not related). | | | | | | | | | | | | | | |



| | | | | | | | | | | | | | |
|---|------------------------------------|---------------------------------------|----|---------------|---|---------------|----|---------------|----|---------------|----------|----|----|
| Name, personal identifier | | Paclitaxel (mg/m ²), A +C | | Starting dose | | Dose-Level -1 | | Dose-Level -2 | | Dose Level -3 | | | |
| | | | | 80 | | 65 | | 50 | | Discontinue | | | |
| | | Carboplatin (AUC), C | | 5 | | 4 | | 3 | | Discontinue | | | |
| Height (cm): | Weight (kg): | Surface (m ²): | | | Reduce starting dose-level in the presence of severe co-morbidity | | | | | | | | |
| Year <input type="text"/> | Week no. | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 |
| | Day, month | | | | | | | | | | | | |
| | | 5. cycle | | | 6. cycle | | | 7. cycle | | | 8. cycle | | |
| Paclitaxel (mg), iv. day 1+8+15 | | | | | | | | | | | | | |
| Carboplatin (mg), iv. day 1 | | | | | | | | | | | | | |
| Growth factor support 0=no, 1=yes | | | | | | | | | | | | | |
| Pembrolizumab (mg) | | | | | | | | | | | | | |
| Ovarian function suppression 0=no, 1=yes | | | | | | | | | | | | | |
| Zoledronic acid (mg) | | | | | | | | | | | | | |
| Hematological toxicity If absolute neutrophil count (ANC) < 1.0 x 10 ⁹ /L on a day of planned treatment postpone treatment until ANC ≥1.0 and add growth factor support. On the second occasion reduce by one level. If platelets < 100 x 10 ⁹ /L on a day of planned treatment postpone treatment until platelets ≥100 and reduce by one dose level. | | | | | | | | | | | | | |
| Enter on day 1 | Neutrophils (x10 ⁹ /L) | | | | | | | | | | | | |
| | Leucocytes (x10 ⁹ /L) | | | | | | | | | | | | |
| | Hemoglobin (mmol/L) | | | | | | | | | | | | |
| | Thrombocytes (x10 ⁹ /L) | | | | | | | | | | | | |
| Renal function | | | | | | | | | | | | | |
| On day 1 | GFR ml/min | | | | | | | | | | | | |
| Non-hematological toxicity Grade 3 and 4 should result in dose reduction | | | | | | | | | | | | | |
| Febrile neutropenia | 0 – 4 | | | | | | | | | | | | |
| Mouth or throat sores | 0 – 2 | | | | | | | | | | | | |
| Nausea | 0 – 2 | | | | | | | | | | | | |
| Vomiting | 0 – 4 | | | | | | | | | | | | |
| Diarrhea | 0 – 4 | | | | | | | | | | | | |
| Rash | 0 – 2 | | | | | | | | | | | | |
| Hand-foot syndrome | 0 – 2 | | | | | | | | | | | | |
| Myalgia | 0 – 2 | | | | | | | | | | | | |
| Arthralgia | 0 – 2 | | | | | | | | | | | | |
| Numbness & tingling | 0 – 4 | | | | | | | | | | | | |
| Cooling gloves used | 0=no, 1=yes | | | | | | | | | | | | |
| Fatigue | 0 – 2 | | | | | | | | | | | | |
| Missed expected menses | 0=no, 1=yes | | | | | | | | | | | | |
| Other, add text | 0 – 4 | | | | | | | | | | | | |
| Other, add text | 0 – 4 | | | | | | | | | | | | |
| In case of any grade 3 or 4 toxicity make a record in the study specific CRF or eJournal of day of start and resolution as well as relation to the study drugs (related, possible related, or not related). | | | | | | | | | | | | | |